



UNIVERSITI PUTRA MALAYSIA

**ELUCIDATION OF THE WNT & AKT/PHOSPHOINOSITIDE-3-KINASE
PATHWAYS IN COLORECTAL CARCINOMA**

KHOR TIN OO

FPSK(P) 2004 2

**ELUCIDATION OF THE WNT & AKT/PHOSPHOINOSITIDE-3-KINASE
PATHWAYS IN COLORECTAL CARCINOMA**

By

KHOR TIN OO

Thesis submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Degree of Doctor of Philosophy

April 2004



Specially dedicated to,

My beloved wife, son (Hong Ze), parents and sister

The memory of,

My grandma and mother-in-law

For their invaluable love, understanding, patience, support and constant faith.

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**ELUCIDATION OF THE WNT & AKT/PHOSPHOINOSITIDE-3-KINASE
PATHWAYS IN COLORECTAL CARCINOMA**

By

KHOR TIN OO

April 2004

Chairman: Professor Seow Heng Fong, Ph.D.

Faculty: Medicine and Health Sciences

Colorectal cancer (CRC) is the third most common cancer in Malaysia and is currently the commonest cancer in males. Genetics, experimental and epidemiological data suggest that CRC develops from complex interaction between inherited susceptibility and environmental factors. Accumulating evidence suggests that the Wnt and PI3K (phosphoinositide-3-kinase)/Akt signalling pathways play a causative role in tumorigenesis of colorectal cancer.

By employing immunohistochemical method, the expression and correlation of several key regulators or related biomolecules of the Wnt and PI3K/Akt signalling pathways in 47 archival formalin fixed, paraffin embedded tissues of surgically resected colorectal cancer (CRC) specimens performed at Kuala Lumpur Hospital (KLH) between 1999 and 2000, were studied. Laser captured microdissection

technique, polymerase chain reaction and direct sequencing were used to investigate mutations in exon 3 of the β -catenin gene. Mutations in the mutation cluster region (MCR) of adenomatous polyposis coli (APC) gene were also investigated. The expressions of Wnt-1, WISP-1 and FRAT-1 mRNA were determined by reverse-transcription and real-time polymerase chain reaction method.

The results showed that: The expressions of Wnt-1, FRAT-1, APC, nuclear β -catenin, cytoplasmic β -catenin, membrane β -catenin, membrane E-cadherin, cytoplasmic E-cadherin, WISP-1, cyclin-D1, p-Akt1 (Ser473), p-Akt1/2/3 (Thr308), p-BAD (Ser136), p-GSK 3 β (Ser9) and survivin were found in 55.3%, 36.2%, 51.1% 44.6%, 95.7%, 30.6%, 46.8%, 95.7%, 31.9%, 10.6%, 34%, 44.7%, 57.4% 44.7% and 59.6% of CRC tissues, respectively and 17.5%, 5% 100%, 0%, 75%, 100%, 100%, 50%, 12.5%, 0%, 5%, 12.5%, 22.5%, 22.5% and 32.5% of apparently normal adjacent tissues, respectively. The sum of scores for all biomolecules except APC, membrane β -catenin and membrane E-cadherin staining was significantly higher in CRC tissues in comparison to apparently normal adjacent tissues ($p < 0.05$). The sum of score for APC, membrane β -catenin and membrane E-cadherin staining was significantly lower in CRC tissues in comparison to apparently normal adjacent tissues ($p < 0.05$). The expression of Wnt and PI3K/Akt signalling pathway-related biomolecules was interrelated. The results of nucleotide sequencing showed that no mutations at exon-3 of β -catenin were found. However, point mutations in the mutation cluster region of the APC gene leading to the formation of truncated APC protein, were found in four

out eleven CRC tissues examined. A 1.43 to 21.26-fold and 1.11 to 109.14-fold increase in the level of expression of Wnt-1 and FRAT-1 mRNA was found in eight out of eleven CRC tissues relative to apparently normal adjacent tissues. On the other hand, a 1.94 to 46.69-fold increase in the level of WISP-1 mRNA was found in all the CRC tissues.

This study has provided important information for researchers and clinicians in terms of clinical evidence of the involvement of the Wnt signalling pathway and PI3K/Akt signalling pathway in colorectal tumorigenesis. In addition, the present study also provided crucial information on the elucidation of the relationship between the biomolecules of these signalling pathways towards understanding their roles in colorectal tumorigenesis and the identification of potential targets for advance therapeutic intervention of CRC. Based on our current results, we propose that Wnt-1, FRAT-1 and WISP-1 could be served as potent therapeutic target for the treatment of CRC.

On the basis of our present study, we conclude that the Wnt and PI3K/Akt signalling pathways are involved in tumorigenesis of CRC in Malaysia. These pathways are interrelated although they might also act independently in promoting tumour growth and inhibition of apoptosis. This study has also provided useful information for the search or design of better antitumour interventions.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PENJELASAN LINTASAN WNT DAN
AKT/PHOSPHOINOSITIDE-3-KINASE DALAM KARSINOMA
KOLOREKTAL**

Oleh

KHOR TIN OO

April 2004

Pengerusi: Profesor Dr Seow Heng Fong, Ph.D.

Fakulti: Perubatan dan Sains Kesihatan

Barah kolorektal (CRC) merupakan barah yang ketiga paling kerap di Malaysia dan pada ketika ini, ia merupakan barah yang paling kerap di kalangan lelaki. Data genetik, eksperimental dan data epidemiologi menyarankan bahawa CRC berkembang hasil interaksi antara faktor persekitaran dan faktor keturunan. Bukti-bukti telah menyarankan bahawa lintasan isyarat PI3K (phosphoinositide-3-kinase) /Akt dan Wnt memainkan peranan yang penting dalam perkembangan barah kolorektal.

Dengan menggunakan kaedah immunohistokimia, ekspresi dan hubungan antara beberapa pengawal-atu atau biomolekul yang berkaitan dengan lintasan isyarat



PI3K/Akt dan Wnt telah dikaji dalam 47 sampel tisu yang diperolehi daripada pesakit CRC yang menjalani pembedahan di Hospital Kuala Lumpur antara 1999 dan 2000 dan telah diblokkan di dalam paraffin. Kaedah “laser captured microdissection”, “polymerase chain reaction (PCR)” dan “direct-sequencing” telah digunakan untuk mengkaji mutasi yang berlaku pada ekson 3, gen β -katenin. Mutasi pada “mutation cluster region (MCR)”, gen APC juga dikaji. Ekspresi mRNA Wnt-1, WISP-1 dan FRAT-1 juga dikenalpasti dengan menggunakan kaedah “real-time”-PCR.

Keputusan kami menunjukkan bahawa ekspresi Wnt-1, FRAT-1, APC, β -katenin nukleus, β -katenin sitoplasma, β -katenin membran, E-cadherin membran, E-cadherin sitoplasma, WISP-1, cyclin-D1, p-Akt1 (Ser473), p-Akt1/2/3 (Thr308), p-BAD (Ser136), p-GSK 3 β (Ser9) dan survivin telah dikesan di 55.3%, 36.2%, 51.1% 44.6%, 95.7%, 30.6%, 46.8%, 95.7%, 31.9%, 10.6%, 34%, 44.7%, 57.4% 44.7% dan 59.6% tisu CRC, masing-masing dan 17.5%, 5% 100%, 0%, 75%, 100%, 100%, 50%, 12.5%, 0%, 5%, 12.5%, 22.5%, 22.5% dan 32.5% tisu sekeliling yang kelihatan biasa, masing-masing. Jumlah skor untuk semua biomolekul kecuali APC, β -katenin membran and E-cadherin membran adalah lebih tinggi dalam tisu CRC berbanding dengan tisu sekeliling yang kelihatan biasa ($p < 0.05$). Jumlah skor untuk APC, β -katenin membran and E-cadherin membran adalah lebih rendah dalam tisu CRC berbanding tisu sekeliling yang kelihatan biasa ($p < 0.05$). Ekspresi biomolekul yang berkaitan dengan lintasan isyarat Wnt dan PI3K/Akt adalah saling berhubungan. Keputusan penjujukan menunjukkan bahawa tidak ada mutasi berlaku di ekson-3,

gen β -katenin. Walau bagaimanapun, mutasi pada MCR, gen APC yang menyebabkan pembentukan protein APC yang “truncated” telah dikesan pada empat daripada sebelas spesimen CRC yang dikaji. Peningkatan ekspresi mRNA Wnt-1 dan FRAT-1 sebanyak 1.43 ke 21.26-kali dan 1.11 ke 109.14-kali masing-masing telah didapati dalam lapan daripada sebelas tisu CRC berbanding tisu sekeliling yang kelihatan biasa. Sebaliknya, peningkatan sebanyak 1.94 ke 46.69-kali mRNA WISP-1 telah berjaya dikesan di kesemua tisu CRC yang dikaji.

Kajian ini telah menghasilkan maklumat yang penting kepada para penyelidik dan perubatan dari segi bukti klinikal bagi pembabitan lintasan isyarat Wnt dan PI3K/Akt dalam tumorigenesis kolorektal. Kajian kami juga memberi maklumat penting dalam penjelasan hubungan antara biomolekul bagi lintasan isyarat yang berkenaan, menuju pemahaman peranan mereka di dalam tumorigenesis kolorektal dan pengenalan sasaran-potensi bagi intervensi CRC terapi maju. Berdasarkan keputusan ini, kami mencadangkan bahawa Wnt-1, FRAT-1 dan WISP-1 boleh dianggap sebagai sasaran terapeutik yang berpotensi untuk rawatan CRC.

Berdasarkan keputusan yang diperolehi, kami membuat kesimpulan bahawa lintasan isyarat Wnt dan PI3K/Akt adalah berkait dengan tumorigenesis CRC di Malaysia. Lintasan isyarat ini adalah saling berhubungan walaupun mereka juga boleh bertindak secara bersendirian untuk menggalakkan pertumbuhan barah dan perencatan apoptosis. Kajian ini telah memberi maklumat yang berguna kepada para

penyelidik dan doctor perubatan dalam penemuan dan penerokaan intervensi anti-barah yang lebih baik.

ACKNOWLEDGEMENTS

I would like to express my sincere gratitude and heartfelt appreciation to my main supervisor, Prof. Dr. Seow Heng Fong, for her advice, patience, encouragement and generous guidance throughout my research. Her supportive supervision and constructive criticism have been crucially important for this thesis.

My sincere thanks are also accorded to my co-supervisors, Prof. Dr. Yunus Gul Alif Gul and Dr. Hairuszah Ithnin, for their endless help, priceless comments and invaluable advice throughout the entire progress of this project.

I would like to acknowledge and thank surgeons Mr. Faisal, Mr. Kwan and Ms. Subhita for generously helping me to collect the specimens. A special note of appreciation also goes out to Dr. Goh Yong Meng for providing logistic support and advice on statistics.

I am indebted to my labmates, Andrea Lisa Holmes, Wang Suk Mei, Maha Abdullah, Ban KeChen, Tina Ong, Leong Pooi Pooi, Lim Pei Ching, Loh Hui Woon and Anthonysamy for their helpful collaboration and discussion.

Finally, I would like to express my deepest gratitude to my wife, parents and sister for their continual support and encouragement for me to proceed to completion of this project.



I certify that an Examination Committee met on 14th April 2004 to conduct the final examination of Khor Tin Oo on his Doctor of Philosophy thesis entitled “Elucidation of the Wnt and Akt/Phosphoinositide-3-Kinase Pathways in Colorectal Carcinoma” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Yushak Abdul Wahab, KMN, SMT. MBBS, FRCS, AM

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Zarida Hambali, MD, DCP, Ph.D., AM

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Mariana Nor Shamsudin, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Alistair Renwick, MA, MBChB, MDPH.D., FRCPATH, FRCPE, CChem, FRCS

Professor
International Medical University
Sesama Street, Plaza Komanwel
Bukit Jalil
57000 Kuala Lumpur
(Independent Examiner)



GULAM RUSUL RAHMAT ALI, Ph.D.
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 30 AUG 2004

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The member of the Supervisory Committee are as follows:

SEOW HENG FONG, Ph.D.

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

YUNUS GUL ALIF GUL, MBBCH, FRCS

Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

HAIRUSZAH ITHNIN, MD, M.PATH, AM

Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.


Professor/Dean,
School of Graduate Studies
Universiti Putra Malaysia

Date: 10 SEP 2004



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



KHOR TIN OO

Date: 15/07/04

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	x
APPROVAL	xii
DECLARATION	xiii
LIST OF TABLES	xx
LIST OF FIGURES	xxii
LIST OF ABBREVIATIONS	xxiv
 CHAPTER	
 I INTRODUCTION	 1-10
 II LITERATURE REVIEW	
2.1 Colorectal carcinoma-General demographics	11
2.2 Risk factors of CRC	12
2.2.1 Host factors	12
2.2.1.1 Longitudinal Cohorts	12
2.2.1.2 Race and ethnicity	13
2.2.1.3 Personal history of colorectal neoplasia	13
2.2.1.4 Insulin and insulin-like growth factor	14
2.2.1.5 Cholecystectomy	15
2.2.1.6 Ulcerative colitis	15
2.2.2 Life-style factors	16
2.2.2.1 Immigration	16
2.2.2.2 Dietary factor	16
2.2.2.3 Physical activity	17
2.3 Clinical manifestation	17
2.4 Diagnosis	18
2.5 Staging	18
2.6 Prognosis	18
2.7 Prevention	22
2.8 Treatment	23
2.8.1 Surgery	23
2.8.2 Chemotherapy	23
2.8.3 Radiotherapy	24
2.8.4 New strategies for CRC treatment	24
2.9 Vogelstein model colorectal tumorigenesis	25
2.10 Wnt signal transduction pathway	27
2.11 Components and alterations of the Wnt signaling pathway in human	30



cancer	
2.11.1 Wnt	30
2.11.2 Frizzled	36
2.11.3 LRP	36
2.11.4 Frizzled related protein (FrzB), Wnt-inhibitory factor-1 (WIF-1) and Dickkopf (Dkk)	38
2.11.5 Dishevelled	39
2.11.6 β -catenin	41
2.11.7 APC	44
2.11.8 Axin	49
2.11.9 GSK-3 β	50
2.11.10 Frat/GBP	52
2.11.11 TCF/LEF	52
2.11.12 Target genes	54
2.11.12.1 Cyclin-D1	54
2.11.12.2 Survivin	54
2.11.12.3 WISP-1	55
2.12 TP53	56
2.13 K-ras	56
2.14 Deleted in colorectal cancer (DCC)	57
2.15 Phosphatidylinositol-3 kinase (PI3-Kinase)/Akt-signaling pathway	57
2.15.1 Involvement of the PI3K/Akt pathway in neoplasia	59
2.15.2 Phosphorylation targets of Akt/PKB	60
2.15.2.1 BAD	61

III MATERIALS AND METHODS 62

3.1 Tissue specimens for immunohistochemistry and laser capture microdissection	63
3.2 Tissue specimens for immunohistochemistry, realtime PCR and PCR	64
3.3 Immunohistochemistry	64
3.3.1 Preparation of sections	64
3.3.2 Standard procedures for immunohistochemical staining	64
3.3.2.1 Immunohistochemical staining of Wnt-1 protein	65
3.3.2.2 Immunohistochemical staining of β -catenin protein	66
3.3.2.3 Immunohistochemical staining of APC protein	66
3.3.2.4 Immunohistochemical staining of E-cadherin protein	66
3.3.2.5 Immunohistochemical staining of FRAT-1 protein	67
3.3.2.6 Immunohistochemical staining of WISP-1 protein	67
3.3.2.7 Immunohistochemical staining of cyclin-D1 protein	67
3.3.2.8 Immunohistochemical staining of survivin protein	68
3.3.2.9 Immunohistochemical staining of phospho-Akt-1 (Ser473) protein	68
3.3.2.10 Immunohistochemical staining of phospho-Akt1/2/3 (Thr308) protein	68
3.3.2.11 Immunohistochemical staining of phospho-GSK-3 β (Ser9) protein	69
3.3.2.12 Immunohistochemical staining of phospho-BAD	69



(Ser136) protein	
3.3.3 Evaluation of immunohistochemical staining	70
3.2.4 Statistical analysis	70
3.4 Detection of gene mutation using laser capture microdissection (LCM)	72
3.4.1 Immunohistochemical staining of β -catenin for LCM	72
3.4.2 Laser capture microdissection	73
3.5 Mutation analysis	73
3.5.1 Extraction of genomic DNA from LCM captured cells and fresh tissues	73
3.5.2 Amplification of the β -catenin gene	74
3.5.3 Amplification of the APC gene	74
3.5.4 Purification and sequencing of PCR products	75
3.5.5 Extraction of total RNA from fresh tissues	76
3.5.6 Semi-quantitative reverse transcription-PCR	77
3.5.6.1 Reverse transcription	77
3.5.6.2 Amplification of Wnt-1 mRNA	78
3.5.6.3 Amplification of WISP-1 mRNA	78
3.5.6.4 Amplification of FRAT-1 mRNA	79
3.5.6.5 Negative and internal control	79
3.5.6.6 Quantification	80
3.6 Real-time RT-PCR	80
3.6.1 Theoretical basis	80
3.6.2 Amplification of Wnt-1, WISP-1 and FRAT-1 mRNA by RTPCR	80
3.6.3 Relative quantification of gene expression using comparative C_T method	81
 IV THE ELUCIDATION OF THE WNT SIGNALING PATHWAY TOWARDS UNDERSTANDING OF ITS ROLE IN COLORECTAL TUMORIGENESIS	 82
4.1 Introduction	82
4.2 Results of the aberrant expression of the Wnt signaling pathway-related biomolecules	83
4.2.1 Detection of Wnt-1 expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	84
4.2.2 Detection of β -catenin expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	86
4.2.3 Detection of mutation in exon-3 of β -catenin genes	87
4.2.4 Detection of APC expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	90
4.2.5 Detection of mutations in mutation cluster region (MCR) of APC genes	91
4.2.6 Detection of E-cadherin expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	94
4.2.7 Detection of FRAT-1 expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	95
4.2.8 Detection of WISP-1 expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	96



4.2.9	Detection of cyclin-D1 expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	97
4.2.10	Detection of survivin expression in apparently normal adjacent tissues and CRC tissues by immunohistochemistry	98
4.2.11	Detection of WISP-1, Wnt-1 and FRAT-1 expression in apparently normal adjacent tissues and CRC tissues by semi-quantitative, conventional RT-PCR and real-time quantitative PCR	99
4.2.11.1	Isolation of total RNA from fresh tissues and reverse transcription	99
4.2.11.2	Amplification of Wnt-1, WISP-1 and FRAT-1 by conventional PCR method	99
4.2.11.3	Semi-quantification of Wnt-1, WISP-1 and FRAT-1	101
4.2.11.4	Real-time amplification of Wnt-1, WISP-1 and FRAT-1 mRNA	102
4.2.11.5	Relative quantification of mRNA expression using comparative C _T method	105
4.3	Discussion on the aberrant expression of the Wnt signaling pathway related biomolecules	106
4.3.1	Wnt-1	106
4.3.2	FRAT-1	108
4.3.3	Adenomatous Polyposis Coli (APC)	109
4.3.4	β -catenin	112
4.3.5	E-cadherin	116
4.3.6	WISP-1	118
4.3.7	Cyclin-D1	119
4.3.8	Survivin	121
4.4	Conclusion on the aberrant expression of the Wnt signaling pathway-related biomolecules	122
4.5	Results of the correlation among the expression of Wnt signaling pathway-related biomolecules	123
4.5.1	Correlation between aberrant expression of β -catenin with other Wnt signaling-related biomolecules	125
4.5.1.1	Correlation between aberrant expression of β -catenin and its adhesion complex counterpart, E-cadherin	125
4.5.1.2	Correlation between aberrant expression of β -catenin and APC	127
4.5.1.3	Correlation between aberrant expression of β -catenin and FRAT-1	128
4.5.2	Elucidation of the role plays by Wnt-1 in the regulation of the Wnt signaling pathway	128
4.5.2.1	Correlation between expression of Wnt-1 and FRAT-1	129
4.5.2.2	Correlation between expression of Wnt-1 and cytoplasmic β -catenin/E-cadherin	129
4.5.2.3	Correlation between expression of Wnt-1 and WISP-1/cyclin-D1	130
4.5.3	Role of E-cadherin and APC as negative regulator of the Wnt-signalling pathway	131
4.6	Conclusion on the correlation among the expression of Wnt signaling pathway-related biomolecules	132

4.7	Results of the association between total biomolecule score with clinicopathological data	134
4.7.1	Association between total biomolecule score with age, gender and race	134
4.7.2	Association between total biomolecule score with histological grade, pathological stage and tumour side	134
4.8	Discussions on the association between total biomolecule score with clinicopathological data	142
4.8.1	Association between total biomolecule score with age, gender and race	142
4.8.2	Association between total biomolecule score with histological grade, pathological stage and tumour side	142
4.9	Conclusions on the association between total biomolecule score with clinicopathological data	144
V	ELUCIDATION OF THE RELATIONSHIP BETWEEN THE BIOMOLECULES OF PHOSPHOINOSITIDE-3 KINASE (PI3K)/AKT SIGNALING PATHWAY IN COLORECTAL CARCINOMA	145
5.1	Introduction	145
5.2	Results of the aberrant expression of PI3K/Akt signaling pathway-related biomolecules	145
5.2.1	Expression of p-Akt1 (Ser ⁴⁷³) in apparently normal adjacent tissues and CRC tissues	146
5.2.2	Expression of p-Akt1/2/3(Thr ³⁰⁸) in apparently normal adjacent tissues and CRC tissues	149
5.2.3	Expression of p-GSK-3 β (Ser ⁹) in apparently normal adjacent tissues and CRC tissues	150
5.2.4	Expression of p-BAD (Ser ¹³⁶) in apparently normal adjacent tissues and CRC tissues	151
5.3	Discussion on the aberrant expression of PI3K/Akt signaling pathway-related biomolecules	152
5.3.1	Expression of Phospho-Akt1/2/3 (Thr308) and phosphor-Akt1(Ser473): indication of the activation of PI3K/Akt signaling pathway	152
5.3.2	Detection of PI3K/Akt signaling pathway-targeted biomolecules	154
5.3.2.1	Phospho-GSK-3 β	154
5.3.2.2	Phospho-BAD (Ser136)	156
5.4	Conclusion on the aberrant expression of PI3K/Akt signaling pathway-related biomolecules	158
5.5	Results of the elucidation of relationship between the biomolecules of PI3K/Akt signaling pathway	158
5.5.1	Correlation between expression of p-Akt1/2/3 (Thr308), p-Akt1 (Ser473) and their downstream substrate p-BAD (Ser136) and p-GSK 3 β (Ser9)	158
5.6	Discussion on the elucidation of relationship between the biomolecules of PI3K/Akt signaling pathway	159



5.7	Conclusion on the elucidation of relationship between the biomolecules of PI3K/Akt signaling pathway	161
5.8	Results of the association between total biomolecule score with clinicopathological data	162
5.9	Discussion on the association between total biomolecule score with clinicopathological data	162
		167
VI POSSIBLE CROSS-TALK BETWEEN THE WNT SIGNALING PATHWAY AND PHOSPHOINOSITIDE-3 KINASE (PI3K)/AKT SIGNALING PATHWAY		
6.1	Introduction	167
6.2	Results of the correlation among the expression of biomolecules	167
6.3	Discussion on the correlation among the expression of biomolecules	169
6.4	Conclusion	172
VII OVERALL CONCLUSION AND RECOMMENDATIONS		173
7.1	Clinical evidence of the involvement of the Wnt and Pi3K/Akt signaling pathway in colorectal tumourigenesis	173
7.2	The elucidation of the relationship between biomolecules in the Wnt signaling pathway and PI3K/Akt signaling pathway	174
7.3	Identification of potential targets for advance therapeutic intervention of CRC	176
7.4	Recommendation	177
REFERENCES		180
APPENDIX		219
BIODATA OF THE AUTHOR		225
MANUSCRIPTS AND ABSTRACT PUBLISHED		

LIST OF TABLES

No.		Page
2.1	Classification of colorectal cancer	19
2.2	TNM staging system for colorectal cancer	20
2.3	Prognostic markers in colorectal cancer	21
2.4	Gene nomenclature of key Wnt components	28
2.5	Overall of mutations of the Wnt signaling pathway in selected human cancer	31
2.6	Mutations in exon 3 of β -catenin (CTNNB1) in various types of cancers	45
3.1	Clinicopathological data of patients	63
3.2	Semi-quantitative scoring system for immunohistochemical staining	71
4.1	Immunohistochemical staining for the expression of various biomolecules	85
4.2	Sum of score for the expression of various biomolecules in apparently normal adjacent tissues and CRC tissues	85
4.3	APC gene mutations of CRC	93
4.4	Correlation among the total scores of biomolecules	124
4.5	Correlation between age and total scores of biomolecules in apparently normal adjacent tissues	135
4.6	Correlation between age and total scores of biomolecules in CRC tissues	136
4.7	Correlation between gender and total scores of biomolecules in apparently normal adjacent tissues	137
4.8	Correlation between gender and total scores of biomolecules in CRC tissues	138
4.9	Correlation between race and total scores of biomolecules in apparently normal adjacent tissues	139

4.10	Correlation between race and total scores of biomolecules in CRC tissues	139
4.11	Correlation between histograde and total scores of biomolecules in CRC tissues	140
4.12	Correlation between pathological stage and total scores of biomolecules in CRC tissues	140
4.13	Correlation between tumor side and total scores of biomolecules in CRC tissues	141
5.1	Immunohistochemical staining for the expression of various biomolecules	147
5.2	Sum of score for the expression of various biomolecules	147
5.3	Correlation among the total scores of biomolecules	159
5.4	Correlation between age and total scores of biomolecules in apparently normal adjacent tissues	163
5.5	Correlation between age and total scores of biomolecules in CRC tissues	163
5.6	Correlation between gender and total scores of biomolecules in apparently normal adjacent tissues	164
5.7	Correlation between gender and total scores of biomolecules in CRC tissues	164
5.8.	Correlation between race and total scores of biomolecules in apparently normal adjacent tissues	165
5.9	Correlation between race and total scores of biomolecules in CRC tissues	165
5.10	Correlation between histograde and total scores of biomolecules in CRC tissues	166
5.11	Correlation between pathological stage and total scores of biomolecules in CRC tissues	166
5.12	Correlation between tumor side and total scores of biomolecules in CRC tissues	166
6.1	Correlation among the total scores of biomolecules	168

LIST OF FIGURES

No.		Page
1.1	A schematic representation of the Wnt signalling pathway	4
1.2	A schematic representation of the PI3K/Akt pathway	8
2.1	Overall scheme of key genetic events in colorectal tumorigenesis	26
2.2	General structure and interaction motif of some important components in the Wnt signaling pathway	37
4.1	Representative slides showing the immunohistochemical staining of Wnt-1	84
4.2	Representative slides showing the immunohistochemical staining of β -catenin	87
4.3	Cells from CRC tissues that expressed nuclear β -catenin were captured by using the laser capture microdissection (LCM) method.	88
4.4	Electropherogram of β -catenin sequence	89
4.5	Representative slides showing the immunohistochemical staining of APC	90
4.6	Amplification of APC gene	91
4.7	Electropherogram of APC sequence	92
4.8	Representative slides showing the immunohistochemical staining of E-cadherin	94
4.9	Representative slides showing the immunohistochemical staining of FRAT-1	95
4.10	Representative slides showing the immunohistochemical staining of WISP-1	96
4.11	Representative slides showing the immunohistochemical staining of cyclin-D1	97

4.12	Representative slides showing the immunohistochemical staining of survivin	98
4.13	The purity of RNA isolated from the fresh CRC and adjacent normal tissues	99
4.14	Amplification of WISP-1 mRNA	100
4.15	Amplification of Wnt-1 mRNA	100
4.16	Amplification of FRAT-1 mRNA	100
4.17	Amplification of 18S gene	101
4.18	Semi-quantitative method for comparison of Wnt-1, FRAT-1 and WISP-1 mRNA in CRC vs normal adjacent tissues	102
4.19	Representative real-time PCR cycling profile of Wnt-1, FRAT-1 and WISP-1 amplification.	103
4.20	Representative melting curve for Wnt-1, FRAT-1 and WISP-1 gene.	104
4.21	Relative quantification of Wnt-1, FRAT-1 and WISP-1.	105
4.22	The outlines of results from (A) previous studies in comparison to the (B) present study on the relationship between the biomolecules in the Wnt signalling pathway	133
5.1	Representative slides showing the immunohistochemical staining of p-Akt1 (Ser473)	148
5.2	Representative slides showing the immunohistochemical staining of p-Akt1/2/3 (Thr308)	149
5.3	Representative slides showing the immunohistochemical staining of p-GSK 3 β (Ser9)	150
5.4	Representative slides showing the immunohistochemical staining of p-BAD (Ser136)	151
5.5	The outlines of results from (A) previous studies in comparison to the (B) present study on the relationship between the biomolecules in the PI3K/Akt signalling pathway	161
6.1	The cross-talk between the Wnt signalling pathway and PI3K/Akt signalling pathway	172

LIST OF ABBREVIATIONS

µg	Microgram
µl	Microlitre
5-FU	5-Fluorouracil
AAPC	Attenuated adenomatous polyposis coli
APC	Adenomatous polyposis coli
bp	Base pair
cAMP	Cyclic adenosine monophosphate
CKIε	Casein kinase I epsilon
Cox	Cyclooxygenase
CRC	Colorectal carcinoma/cancer
DCC	Deleted in colorectal cancer
Dkk	Dickkopf
DNA	Deoxyribonucleic acid
dNTPs	Dideoxynucleotide triphosphates
Dsh/Dvl	Dishevelled
EGF	Epidermal growth factor
FAP	Familial adenomatous polyposis
FOBT	Faecal occult blood test
FRAT	Frequently rearranged in advance T-cell lymphocytes
FrzB	Frizzled-related protein
Fz	Frizzled
g	Gram
GBP	Glycogen synthase kinase binding protein
GSK3β	Glycogen synthase kinase 3β
HNPCC	Hereditary non-polyposis colorectal cancer
IAP	Inhibitor of apoptosis protein
IGF	Insulin-like growth factor
JNK	Jun kinase
kb	Kilobase pair
LCM	Laser capture microdissection
LEF	Lymphoid enhancer factor
LRP	Low density lipoprotein-receptor-related protein
mA	Milliampere
mg	Milligram
MgCl ₂	Magnesium Chloride
mRNA	messenger Ribonucleic acid
MUC1	Mucin antigen 1
nM	Nano molar
NHS	Nurses' health study
Nkd	Naked cuticle
p-Akt	Phosphorylated Akt
p-BAD	Phosphorylated BAD
PBS	Phosphate buffer saline